

Answers to Selected Exercises in *Real and Complex Analysis**

Solutions by

MARCEL K. GOH

CHAPTER 1. ABSTRACT INTEGRATION

2. Put $f_n = \chi_E$ if n is odd, $f_n = 1 - \chi_E$ if n is even. What is the relevance of this example to Fatou's lemma?

This gives an example of strict inequality arising. Let X be a measure space such that $\mu(X) = 1$ and let $E \subseteq X$ be such that $\mu(E) = 2/3$. Then

$$\int_X \liminf_{n \rightarrow \infty} f_n d\mu = \int_X 0 d\mu = 0,$$

while on the other hand

$$\liminf_{n \rightarrow \infty} \int_X f_n d\mu = \int_X 1 - \chi_E d\mu = \int_X 1 d\mu - \int_X \chi_E d\mu = \mu(X) - \mu(E) = \frac{1}{3},$$

and we have $0 < 1/3$.

3. Suppose $f_n : X \rightarrow [0, \infty]$ is measurable for $n = 1, 2, 3, \dots$, $f_1 \geq f_2 \geq f_3 \geq \dots \geq 0$, $f_n(x) \rightarrow f(x)$ as $n \rightarrow \infty$ for every $x \in X$, and $f_1 \in L^1(\mu)$. Prove that then

$$\lim_{n \rightarrow \infty} \int_X f_n d\mu = \int_X f d\mu$$

and show that this conclusion does not follow if the condition " $f_1 \in L^1(\mu)$ " is omitted.

Proof. Consider the sequence of functions $(f_1 - f_n)$. This sequence is nonnegative, nondecreasing and for any $x \in X$, $\lim_{n \rightarrow \infty} (f_1 - f_n)(x) = f_1(x) - f(x)$. So by the monotone convergence theorem,

$$= \lim_{n \rightarrow \infty} \int_X f_1 - f_n d\mu = \int_X f_1 - f d\mu.$$

Since the integrals of f_n are finite, we can apply Theorem 1.27 to get

$$\int_X f_1 d\mu - \lim_{n \rightarrow \infty} \int_X f_n d\mu = \int_X f_1 d\mu - \int_X f d\mu,$$

which gives us what we want after subtracting $\int_X f_1 d\mu$ (which is finite) and multiplying by -1 . ■

The condition $f_1 \in L^1(\mu)$ is necessary. Let $I = [0, 1]$ and $f_n = \frac{1}{nx^2}$, which converges to the constant function 0. Then $\lim_{n \rightarrow \infty} \int_I f_n d\mu = +\infty$ but $\int_I 0 d\mu = 0$.

4. Prove that if f is a real function on a measurable space X such that $\{x : f(x) \geq r\}$ is measurable for every rational r , then f is measurable.

Proof. We immediately infer that for rational r and s , $f^{-1}([r, s))$ is measurable, since

$$f^{-1}([r, s)) = \{x : f(x) \geq r\} \cap \{x : f(x) \geq s\}^c.$$

* Walter Rudin. 1966. *Real and complex analysis*. McGraw-Hill, Inc., USA.

Next, let $I \subseteq \mathbf{R}$ be an open interval. Setting $J = I \cap \mathbf{Q}$, we can show that $f^{-1}(I)$ is measurable, since

$$f^{-1}(I) = f^{-1}\left(\bigcup_{r \in J} \bigcup_{s \in J} [r, s)\right) = \bigcup_{r \in J} \bigcup_{s \in J} f^{-1}([r, s))$$

is a countable union of measurable sets.

Now let $V \subseteq \mathbf{R}$ be open. By Lindelöf's lemma, we can express V as a countable union of open intervals $V = \bigcup_{n=1}^{\infty} I_n$. Then

$$f^{-1}(V) = f^{-1}\left(\bigcup_{n=1}^{\infty} I_n\right) = \bigcup_{n=1}^{\infty} f^{-1}(I_n)$$

is measurable. \blacksquare

5. Prove that the set of points at which a sequence of measurable real-valued functions converges (to a finite limit) is measurable.

Proof. Let (f_n) be a sequence of real-valued measurable functions. Then (f_n) converges at a point x if and only if it is Cauchy at x , i.e. for any $\epsilon > 0$ there exists $N \in \mathbf{N}$ such that for all $m, n \geq N$, $|f_m(x) - f_n(x)| < \epsilon$. By the Archimedean property, we can replace ϵ with $1/k$ for some $k \in \mathbf{N}$. So the set of all points at which (f_n) converges can be written thus:

$$\bigcap_{k=1}^{\infty} \bigcup_{N=1}^{\infty} \bigcap_{m=1}^{\infty} \bigcap_{n=1}^{\infty} \{x : |f_m(x) - f_n(x)| < 1/k\}$$

Set $g_{m,n} = f_m - f_n$; this is a measurable function. Hence the set

$$\{x : |f_m(x) - f_n(x)| < 1/k\} = \{x : |g_{m,n}(x)| < 1/k\} = g_{m,n}^{-1}((-1/k, 1/k))$$

is measurable for every m, n , and k . We have thus proved that the set of all points at which (f_n) converges is a countable union of measurable sets. \blacksquare

6. Let X be an uncountable set, let \mathcal{M} be the collection of all sets $E \subseteq X$ such that either E or E^c is at most countable, and define $\mu(E) = 0$ in the first case, $\mu(E) = 1$ in the second. Prove that \mathcal{M} is a σ -algebra in X and that μ is a measure on \mathcal{M} . Describe the corresponding measurable functions and their integrals.

First we prove that \mathcal{M} is a σ -algebra and that μ is a measure.

Proof. Since \emptyset is countable, $X \in \mathcal{M}$. Then for any set $E \in \mathcal{M}$, either E or E^c is at most countable; in any case $E^c \in \mathcal{M}$. Lastly, let $\{E_n\}$ be a collection of sets in \mathcal{M} . If every set E_n is at most countable, then $\bigcup_{n=1}^{\infty} E_n$ is countable and thus in \mathcal{M} . Otherwise, there is some k for which E_k is uncountable. Then $E_k^c \in \mathcal{M}$ is at most countable and $(\bigcup_{n=1}^{\infty} E_n)^c = \bigcap_{n=1}^{\infty} E_n^c \subseteq E_k^c$ is countable, so $\bigcup_{n=1}^{\infty} E_n$ is in \mathcal{M} . This shows that \mathcal{M} is a σ -algebra.

It is clear that the range of μ is in $[0, \infty]$. To show that μ is countably additive, let $\{E_n\}$ be a disjoint collection of sets in \mathcal{M} . If every E_n is at most countable, then $\bigcup_{n=1}^{\infty} E_n$ is at most countable and $\mu(\bigcup_{n=1}^{\infty} E_n) = 0 = \sum_{n=1}^{\infty} \mu(E_n)$. Otherwise, there exists E_k such that E_k^c is countable and thus we have $\mu(E_k) = 1$ and $\mu(E_k^c) = 0$. Since the E_n are pairwise disjoint, $E_n \subseteq E_k^c$ for all $n \neq k$. So $\mu(E_n) = 0$ for all $n \neq k$. So $\mu(\bigcup_{n=1}^{\infty} E_n) = 1 = \sum_{n=1}^{\infty} \mu(E_n)$ and μ is a measure. \blacksquare

Now we claim that the measurable functions are those that are constant at all but countably many points.

Proof. It suffices to prove this for a real-valued function. Suppose $f : X \rightarrow [-\infty, \infty]$ is measurable. For any $a \in \mathbf{R}$, let $E_a = f^{-1}([-\infty, a))$. Note that for any a , either E_a is countable or E_a^c is countable. Note also that if $a \leq b$, then $E_a \subseteq E_b$. So let there is a constant c such that

$$k = \sup\{a : E_a \text{ is countable}\}.$$

This supremum is not $-\infty$, since if it were, then E_a^c would be countable for all $a \in \mathbf{R}$, and since $\bigcap_{n=0}^{\infty} E_{-n} = \emptyset$, $X = \bigcap_{n=0}^{\infty} E_{-n}^c$ is countable, a contradiction. By a similar argument, we know that the supremum is not

∞ . So $k \in \mathbf{R}$ and there exists a sequence (a_n) all of whose terms are less than k and whose limit is k . Hence $E_k = \bigcup_{n=1}^{\infty} E_{a_n}$ is countable. Now let (b_n) be a sequence that converges to k such that $b_n > k$ for all n . Then note that if $f(x) > k$, then $f(x) \in [b_n, \infty) = E_{b_n}^c$ for some n . So

$$\{x : f(x) \neq k\} = E_k \cup \{x : f(x) > k\} \subseteq E_k \cup \left(\bigcup_{n=0}^{\infty} E_{b_n}^c \right)$$

is countable and f equals k at all but countably many points. ■

Lastly, we show that if f is measurable and takes on the value k at all but countably many points, then $\int_E f d\mu = k$ for all uncountable $E \in \mathcal{M}$. (If E is countable then $\int_E f d\mu = 0$.)

Proof. Let $E \in \mathcal{M}$ be uncountable; so $\mu(E) = 1$. Let s be a simple measurable function such that $0 \leq s \leq f$. Suppose that s takes values α_i on the n disjoint sets A_i that cover X . We know that one of the α_i , call it α_j , is equal to a constant k_s and that A_i is at most countable for all $i \neq j$. So for all $i \neq j$, $A_i \cap E$ is countable and $A_j \cap E$ must be uncountable. Then we have

$$\int_E s d\mu = \sum_{i=1}^n \alpha_i \mu(A_i \cap E) = \alpha_j \mu(A_j \cap E) = k_s \cdot 1 = k_s,$$

where $0 \leq k_s \leq k$. So we have

$$\int_E f d\mu = \sup \left\{ \int_E s d\mu : 0 \leq s \leq f \right\} = \sup \{k_s : 0 \leq s \leq f\} = k,$$

which is what we had to show. ■

7. Does there exist an infinite σ -algebra which has only countably many members?

The answer is no.

Proof. Let X be a ground set and let \mathcal{F} be a σ -algebra (over X) with infinitely many members. First we claim that we can always find a set $E \neq \emptyset$ such that the set $\{F \cap E^c : F \in \mathcal{F}\}$ is infinite. If this did not hold, then take any $E \neq \emptyset$ in \mathcal{F} . Then by assumption, $\mathcal{S}_1 = \{F \cap E^c : F \in \mathcal{F}\}$ is finite and because $E^c \in \mathcal{F}$, $\mathcal{S}_2 = \{F \cap E : F \in \mathcal{F}\}$ is finite as well. Since any member of \mathcal{F} can be expressed as a union of an element of \mathcal{S}_1 with an element of \mathcal{S}_2 , this implies that \mathcal{F} is finite, a contradiction.

Now we may use the claim to construct a countable sequence of pairwise disjoint elements of \mathcal{F} . Let G_1 be the set E given by the claim. Now the infinite set \mathcal{S}_1 we constructed before is also a σ -algebra, so repeat the argument to get a set G_2 , disjoint from G_1 . Continuing in this manner, we obtain a sequence G_1, G_2, \dots where the G_i are pairwise disjoint. Now we see that the map from the power set of the natural numbers to \mathcal{F} given by

$$A \mapsto \bigcup_{i \in A} G_i$$

is injective. So the uncountability of \mathcal{F} follows from the uncountability of $\mathcal{P}(\mathbf{N})$. ■